

Summary

SAD detected 99 square kilometers of deforestation in the Brazilian Amazon in November 2015. That represented a 49% reduction in relation to November 2014 when deforestation totaled 195 square kilometers. It was possible to monitor 80% of the forest area in the Brazilian Amazon, while in November 2014 monitoring covered a smaller area (67%) of the territory.

In November 2015, deforestation was concentrated in Mato Grosso (33%), Pará (24%), Rondônia (19%) and Amazonas (19%), with a lower occurrence in Roraima (2%) and Acre (2%).

Degraded forests in the Brazilian Amazon totaled 362 square kilometers in November 2015. In relation to November 2014 when forest degradation totaled 86 square kilometers, there was a 323% increase.

Antônio Fonseca,
Marcelo Justino,
Carlos Souza Jr.
& Adalberto Veríssimo
(Imazon)

Deforestation Statistics

According to SAD, deforestation (total suppression of the forest for other alternative land uses) reached 99 square kilometers in November 2015 (Figure 1 and Figure 2).




Figure 1. Deforestation from August 2014 to November 2015 in the Brazilian Amazon (Source: Imazon/SAD).




Figure 2. Deforestation and Forest Degradation in the Brazilian Amazon in November 2015 (Source: Imazon/ SAD).

In November 2015, deforestation was concentrated in Mato Grosso (33%), Pará (24%), Rondônia (19%) and Amazonas (19%), with a lower occurrence in Roraima (2%) and Acre (2%) (Figure 3).

The deforestation accumulated during the period from August to November 2015, corresponding to the first four months of the official calendar for measuring deforestation, affected 972 square kilometers. There was a 24% reduction in deforestation in relation to the previous period (August 2014 to November 2014) when it reached 1,278 square kilometers.




Figure 3. Percentage of deforestation in States of the Brazilian Amazon in August 2015 (Source: Imazon/SAD).

Evolution of deforestation among States of the Brazilian Amazon in August 2014 and August 2015 (Source: Imazon/SAD).

State	August to November 2014	August to November 2015	Variation (%)
Pará	343	293	-15
Mato Grosso	312	342	+10
Rondônia	327	138	-58
Amazonas	173	133	-23
Roraima	52	29	-44
Acre	66	12	-81
Tocantins	1	25	+2.400
Amapá	4	-	-
Total	1,278	972	-24

* Os dados do Maranhão não foram analisados.

Forest Degradation

In November 2015, SAD recorded 362 square kilometers of degraded forests (forests intensely exploited by timber activity and/or burned) (Figures 2 and 4). Of that total, the majority (49%) occurred in Mato Grosso, followed by Pará (37%) and with lower occurrence in Amazonas (7%) and Rondônia (6%).

Figure 4. Forest Degradation from August 2014 to November 2015 in the Brazilian Amazon (Source: Imazon/SAD).




Table 2. Evolution of forest degradation among the States in the Brazilian Amazon from August 2014 to November 2014 and August 2015 to November 2015 (Source: Imazon/SAD).

State	August to November 2014	August to November 2015	Variation (%)
Mato Grosso	1.153	1.820	+58
Pará	65	258	+297
Rondônia	35	77	+120
Amazonas	7	40	+471
Roraima	4	5	+25
Acre	-	-	-
Tocantins	-	-	-
Amapá	2	-	-100
Total	1,266	2,200	+74

Geography of Deforestation

In November 2015, the majority (61%) of deforestation occurred in areas that were private or under various stages of possession. The remaining deforestation was recorded in Land Reform Settlements (25%), Conservation Units 11% and Indigenous Lands (3%) (Table 3).

Table 3. Deforestation by land title category in November 2015 in the Brazilian Amazon (Source: Imazon/ SAD).

Category	November 2014	
	km ²	%
Land Reform Settlement	25	25
Conservation Units	11	11
Indigenous Lands	3	3
Private, Possession & Untitled Public Lands	60	61
Total (km²)	99	100

Land Reform Settlements

SAD recorded 25 square kilometers of deforestation in Land Reform Settlements in November 2015 (Figure5). The Settlements most affected by deforestation were PAE Novo Tempo Ilha Muratu (Iranduba; Amazonas), PA Nova Cotriguaçu (Cotriguaçu; Mato Grosso) and PA Monte (Boca do Acre; Amazonas).




Figure 5. Most deforested Land Reform Settlements in the Brazilian Amazon in November 2015 (Source: Imazon/SAD).

Protected Areas

In the month of November 2015, SAD detected 11 square kilometers of deforestation in Conservation Units (Figure 6). In the case of Indigenous Lands, in November 2015 3 square kilometers of deforestation were detected (Figure 7).




Figure 6. Most deforested Conservation Units in the Brazilian Amazon in November 2015
(Source: Imazon/SAD).




Figure 7. Most deforested Indigenous Lands in the Brazilian Amazon in November 2015
(Source: Imazon/SAD).

Critical Municipalities

In November 2015, the most deforested municipalities were: Colniza (Mato Grosso) and Manicoré (AM) (Figure 8 and 9).




Figure 8. Most deforested Municipalities in the Brazilian Amazon in November 2015
(Source: Imazon /SAD).




Figure 9. Municipalities com largest areas deforested in the Brazilian Amazon in November 2015
(Source: Imazon/SAD).

Cloud and Shadow Cover

In November 2015, with SAD it was possible to monitor 80% of the forest area in the Brazilian Amazon. The other 20% of forest territory was covered by clouds, which made detection of deforestation and forest degradation difficult. The States with the largest cloud cover were Amapá (70%) and Pará (30%). Because of that, the data for deforestation and forest degradation in November 2015 may be underestimated (Figure 10).




Figure 10. Area with cloud and shadow in November 2015 in the Brazilian Amazon

SAD-EE

Since August 2012 deforestation and forest degradation detection alerts have been performed using the Google Earth Engine (EE), platform (EE), with the new SAD EE version. That system was developed in collaboration with Google and uses the same process already employed by SAD 3.0 (Box I), with reflectance images from MODIS for generating deforestation and forest degradation alerts.

* The part of Maranhão that belongs to the Brazilian Amazon was not analyzed.

Box I: SAD 3.0

Since August 2009, SAD has had some new features. First, we created a graphic interface to integrate all of the image processing programs used in SAD. Next, we began to compute deforestation in areas that were covered by clouds in the previous months in a new class. Finally, deforestation and degradation are detected with pairs of NDFI images using a change detection algorithm. The principal method continues to be the same as with SAD 2.0 as described below.

SAD generates a temporal mosaic of daily MODIS images from the MOD09GQ and MOD09GA products for filtering clouds. Next, we use a technique for fusing different spectral resolution bands, i.e. with pixels of different sizes. In this case, we made a change in scale from 5 bands with 500 meter pixels in MODIS to 250 meters. That allowed us to improve the spectral mixture model and provided the capacity for estimating the abundance of Vegetation, Soils and Non-Photosynthetic Vegetation (NPV) components (Vegetation, Soil and Shadow) to calculate the NDFI, with the following equation:

$$\text{NDFI} = \frac{\text{VGs} - (\text{NPV} + \text{Soil})}{\text{VGs} + \text{NPV} + \text{Soil}}$$

Where VGs is the Vegetation component normalized for shadow given by:

$$\text{VGs} = \text{Vegetation} / (1 - \text{Shadow})$$

The NDFI varies from -1 (pixel with 100% of exposed soil) to 1 (pixel with > 90% of forest vegetation). Thus, we have a continuous image that shows the transition from deforested areas, going through degraded forests, until reaching forest without signs of disturbances.

Detection of deforestation and degradation this month involved a difference in the NDFI images from consecutive months. Thus, a reduction in the NDFI values of from -200 to -50 indicates possible deforested areas and from -49 to -20 indicates signs of degradation.

SAD 3.0 Beta is compatible with previous versions (SAD 1.0 and 2.0), because the threshold for detecting deforestation was calibrated to generate the same type of response obtained by the previous method.

SAD has been in operation in the State of Mato Grosso since August 2006 and in the Brazilian Amazon since May 2008. In this bulletin, we present the monthly data generated by SAD from August 2014 to November 2015.

Team Responsible:

General Coordination: Carlos Souza Jr. and Adalberto Veríssimo (Imazon)
Technical Coordination: Antônio Fonseca
Team: Marcelo Justino and Dalton Cardoso (Image interpretation),
Kátia Pereira and Victor Lins (ImazonGeo), Stefânia Costa (Communications)

Data Source:

Statistics on deforestation are generated using data from SAD (Imazon);
Data from INPE - Deforestation (PRODES)
<http://www.obt.inpe.br/prodes/>

Acknowledgment:

Google Earth Engine
<http://earthengine.google.org/>

Support:

Gordon & Betty Moore Foundation
United States Agency International Development

Partnerships:

State Secretariat for the Environment and Sustainability of Pará (SEMAS)
State Secretariat for the Environment of Mato Grosso (SEMA)
Secretariat for the Environment of Pará (SEMA)
Federal Public Prosecution Service in Pará
State Public Prosecution Service of Pará
State Public Prosecution Service of Roraima
State Public Prosecution Service of Amapá
State Public Prosecution Service of Mato Grosso
Instituto Centro de Vida (ICV- Mato Grosso)

Apoio

GORDON AND BETTY
MOORE
FOUNDATION

